

© 2021 Heldermann Verlag  
Journal of Convex Analysis 28 (2021) 457–470

**F. Cagnetti**

Department of Mathematics, University of Sussex, Brighton, United Kingdom  
f.cagnetti@sussex.ac.uk

**A. Chambolle**

CMAP – CNRS, Ecole Polytechnique, Palaiseau, France  
antonin.chambolle@cmap.polytechnique.fr

**M. Perugini**

Institut für numerische und angewandte Mathematik, Westf.-Wilhelms-Universität, Münster,  
Germany  
matteo.perugini@uni-muenster.de

**L. Scardia**

Department of Mathematics, Heriot-Watt University, Edinburgh, United Kingdom  
l.scardia@hw.ac.uk

**An Extension Result for Generalised Special Functions of Bounded Deformation**

We show an extension result for generalised special functions of bounded deformation ( $GSBD^p$ , for every  $p > 1$ ) and any dimension  $n \geq 2$ . The proof is based on a recent result of F. Cagnetti, A. Chambolle, and L. Scardia, showing that a function  $u$  in  $GSBD^p$  with a “small” jump set coincides with a  $W^{1,p}$  function, up to a small set whose perimeter and volume are controlled by the size of the jump of  $u$ .

**Keywords:** Free-discontinuity problems, functions of bounded deformation, Griffith’s energy.

**MSC:** 49Q20, 70G75, 74R10.