

© 2010 Heldermann Verlag
Journal of Lie Theory 20 (2010) 409–436

F. B. Gonzalez

Dept. of Mathematics, Tufts University, Medford, MA 02155, U.S.A.
fulton.gonzalez@tufts.edu

Conical Distributions on the Space of Flat Horocycles

Let $G_0 = \mathfrak{p} \rtimes K$ be the Cartan motion group associated with a noncompact semisimple Riemannian symmetric pair (G, K) . Let \mathfrak{a} be a maximal abelian subspace of \mathfrak{p} and let $\mathfrak{p} = \mathfrak{a} + \mathfrak{q}$ be the corresponding orthogonal decomposition. A flat horocycle in \mathfrak{p} is a G_0 -translate of \mathfrak{q} . A conical distribution on the space Ξ_0 of flat horocycles is an eigendistribution of the algebra $\mathbb{D}(\Xi_0)$ of G_0 -invariant differential operators on Ξ_0 which is invariant under the left action of the isotropy subgroup of G_0 fixing \mathfrak{q} . We prove that the space of conical distributions belonging to each generic eigenspace of $\mathbb{D}(\Xi_0)$ is one-dimensional, and we classify the set of all conical distributions on Ξ_0 when G/K has rank one. We also consider the question of the irreducibility of the natural representation of G_0 on the eigenspaces of $\mathbb{D}(\Xi_0)$.

Keywords: Conical distributions, Cartan motion group, horocycle Radon transform.

MSC: 43A85; 22E46, 44A12